

Abundances of Carbon Monoxide, Formaldehyde, and Methyl Alcohol in Comets: Measuring Efficiencies for Conversion of CO in Grain Mantles

Michael A. DiSanti

*Goddard Center for Astrobiology
and
Solar System Exploration Division
NASA's Goddard Space Flight Center
Greenbelt, MD 20771
USA
disanti@lepvax.gsfc.nasa.gov*

Michael J. Mumma

*Goddard Center for Astrobiology
and
Solar System Exploration Division
NASA's Goddard Space Flight Center
USA*

Boncho P. Bonev

*University of Toledo at Solar System Exploration Division, NASA's GSFC
USA*

Neil Dello Russo

*Catholic University of America at Solar System Exploration Division, NASA's GSFC
USA*

Karen Magee-Sauer

*Dept. of Chemistry and Physics
Rowan University
Glassboro, NJ
USA*

William M. Anderson

*Catholic University of America at Solar System Exploration Division, NASA's GSFC
USA*

Dennis C. Reuter

*Solar System Exploration Division
NASA's Goddard Space Flight Center
USA*

Erika Gibb

*Dept. of Physics
University of Notre Dame
South Bend, IN
USA*

Hydrogen addition to CO ice in grain mantles prior to their incorporation into comets has been proposed as a viable means of producing monomeric formaldehyde and methanol, a

process that has been verified in laboratory irradiation experiments on pre-cometary analog ices ($\text{H}_2\text{O}:\text{CO}$ mixtures). The relative abundances of CO , H_2CO , and CH_3OH in comets represents one measure of the conversion efficiency of CO , and provides information on extant conditions (e.g., H-atom densities) in the region of the proto-solar nebula where these comets formed. Using high-resolution echelle spectrometers, we have detected CO and CH_3OH in ten Oort cloud comets, and at least four of these also exhibit definitive emission from H_2CO . We will report measured abundances and conversion efficiencies for several comets in our database.

The presence of formaldehyde and related molecules in comets can provide fundamental information pertaining to the origin of life. For example, H_2CO not only is significant in its polymerization to sugar but, along with HCN and NH_3 , is thought to play an important role in the synthesis of amino acids in primitive bodies such as the parent of the Murchison meteorite.

We also apply an existing fluorescence model for H_2CO to line-by-line intensities in our comet spectra. This provides a constraint on the rotational temperature for comparison with that measured for other molecules (e.g., CO , HCN , H_2O), and permits more accurate retrieval of production rates.

This work is supported by grants 344-53-51, 344-33-55, and 344-32-98 from the NASA Astrobiology, Atmospheres, and Astronomy Programs, respectively.